arXiv:astro-ph/0209007v1 31 Aug 2002

X-ray Evidence for Supermassive Black Holes at the Centers of Nearby, Non-Active Galaxies

Stefanie Komossa

Max-Planck-Institut für extraterrestrische Physik
Postfach 1312, 85741 Garching, Germany
skomossa@mpe.mpg.de, http://www.xray.mpe.mpg.de/∼skomossa/

Abstract

We first present a short overview of X-ray probes of the black hole region of active galaxies (AGN) and then concentrate on the X-ray search for supermassive black holes (SMBHs) in optically non-active galaxies.

The first part focuses on recent results from the X-ray observatories Chandra *and* XMM-Newton *which detected a wealth of new spectral features which originate in the nuclear region of AGN.*

In the last few years, giant-amplitude, non-recurrent X-ray flares have been observed from several non-active *galaxies. All of them share similar properties, namely: extreme X-ray softness in outburst, huge peak luminosity (up to* $\sim 10^{44}$ *erg/s), and the absence of optical signs of Seyfert activity. Tidal disruption of a star by a supermassive black hole is the favored explanation of these unusual events. The second part provides a review of the initial X-ray observations, follow-up studies, and the relevant aspects of tidal disruption models studied in the literature.*

1 Introduction

1.1 The search for SMBHs at the centers of galaxies

The study of supermassive black holes and their cosmological evolution is of great interest for a broad range of astrophysical topics including facets of galaxy formation and general relativity. In the last few decades, a number of different methods were developed to search for supermassive black holes (SMBHs) in external galaxies. Their detection in large numbers would clarify our understanding of the early phases of the evolution of galaxies. In *active* galactic nuclei SMBHs are now generally believed to be the prime mover of the non-stellar activity. X-ray observations in the near future are expected to offer the opportunity of detecting some of the distinctive features of strong field gravity, thereby also providing the ultimate proof for the existence of black holes in AGN.

There is now strong evidence for the presence of massive dark objects at the centers of many galaxies. Does this hold for *all* galaxies ? If so, why are some SMBHs 'dark' ? Questions of particular interest in the context of galaxy/AGN evolution are: When and how did the first SMBHs form and how do they evolve ? What fraction of galaxies have passed through an active phase, and how many now have non-accreting and hence unseen supermassive black holes at their centers (e.g., Lynden-Bell 1969, Rees 1989)?

Several approaches were followed to study these questions. Much effort has concentrated on the determination of central object masses from measurements of the *dynamics of stars and gas* in the nuclei of nearby galaxies. Earlier (ground-based) evidence for central quiescent dark masses in galaxies (Kormendy & Richstone 1995) has been strengthened by recent HST results (see Kormendy & Gebhardt 2001 for a review).

A quite accurate determination of black hole mass was enabled by the detection of water vapor maser emission from the mildly active galaxy NGC 4258 (Miyoshi et al. 1995). The water masers, whose motion can be precisely mapped with VLBI, are located in a very compact disk in Keplerian rotation around the central SMBH. The fortunate geometry of the disk, nearly edge-on, allows to obtain the BH mass with high accuracy: $M_{\rm BH} = 3.6\,10^7$ $\mathrm{M}_\odot$ (Neufeld & Maloney 1995, Greenhill et al. 1995).

Still closer to the SMBH, in *active* galaxies with with broad line region (BLR hereafter), the technique of BLR reverberation mapping (e.g., Peterson 2001) provides a powerful tool to estimate the BH mass via the clouds' distance from the center and their velocity field (e.g., Peterson & Wandel 2000, Ferrarese et al. 2001).

1.2 X-ray probes of the black hole region of AGN

Whereas the dynamics of stars and gas probe rather large distances from the SMBH, high-energy *X-ray emission* originates from the immediate vicinity of the black hole. In *active* galaxies, excellent evidence for the presence of SMBHs is provided by the detection of luminous hard power-law like X-ray emission, rapid variability, and the discovery of evidence for relativistic effects in the iron-K line profile. X-ray observations currently provide the most powerful way to explore the black hole region of AGN.

X-rays at the centers of AGN arise in the accretion-disk – corona system (e.g, Mushotzky et al. 1993, Svensson et al. 1994, Collin et al. 2000, and references therein). On larger scales, but still within the central region, X-rays might be emitted by a hot intercloud medium at distances of the broad or narrow-line region (e.g., Elvis et al. 1990).

The X-rays which originate from the accretion-disk region are reprocessed in form of absorption and partial re-emission (e.g., George & Fabian 1991, Netzer 1993, Krolik & Kriss 1995, Collin-Souffrin et al. 1996, Komossa & Fink 1997b) as they make their way out of the nucleus. The reprocessing bears the disadvantage of veiling the *intrinsic* X-ray spectral shape, and the spectral disentanglement of many different potentially contributing components is not always easy. However, reprocessing also offers the unique chance to study the physical conditions and dynamical states of the reprocessing material (see Komossa 2001 for a review), like: the outer parts of the accretion disk; the ionized absorber; the torus, which plays an important role in AGN unification schemes (Antonucci 1993); and the BLR and NLR. Detailed modeling of the reprocessor(s) is also necessary to recover the shape and properties of the *intrinsic* X-ray spectrum.

Figure 1: Sketch of the central region of Seyfert galaxies. The black hole and accretion disk region is surrounded by two systems of gas clouds, the broad line region (BLR) and narrow-line region (NLR). These show up by their characteristic line emission in optical spectra of AGN, and their presence is usually used to identify and classify AGN. The molecular torus, and variants of it, are thought to play an important role in unification schemes of Seyfert galaxies by blocking the direct view on the BLR for certain viewing directions of the observer (Antonucci 1993). Somewhere outside the bulk of the BLR, a relatively recently discovered component of the active nucleus is located, the so-called 'warm' or ionized absorber (WA).
A modification of this picture was recently proposed by Elvis (2000, 2001). In his model, the BLR clouds arise from a flow of gas which rises vertically from a narrow range of radii from the accretion disk. The flow then bends and forms a conical wind moving radially outwards. The BLR clouds are identified with the cool phase of this two-phase medium. Warm absorbers appear if we view the continuum source through the wind (Elvis 2000, see his Fig. 1)

Recent progress has been made based on the improved spectral resolution of the new generation of X-ray observatories, *Chandra* and *XMM-Newton*. Both missions have imaging detectors and grating spectrometers aboard. Their energy sensitivity bandpass covers ~(0.1–10) keV. Below, a short review of results from these observatories is given, starting at relatively large distances from the SMBH (NLR), and then moving further inward (warm absorber and accretion disk region).

X-ray emission lines, X-ray narrow-line region. The detection of a high-temperature, narrow-line, X-ray emitting plasma in NGC 4151 was reported by Ogle et al. (2000), confirming earlier evidence for extended X-ray emission from this galaxy (Elvis et al. 1983). The X-ray gas is spatially coincident with the NLR and extended narrow-line region. For the first time, numerous emission-lines were detected in the X-ray spectrum of NGC 4151 with the HETG (High Energy Transmission Grating Spectrometer) aboard *Chandra*.

The X-ray emission lines detected in the spectrum of NGC 4151 and several Seyfert 2 galaxies (e.g., Mrk 3, NGC 1068), contain important information on the physical conditions in the line-emitting medium, like temperature, density, and the main gas excitation/ionization mechanism - photoionization or collisional ionization. Of particular importance in determining the main power mechanism of the lines, are the Helium-like triplets (Gabriel & Jordan 1969; see our Fig.2), the widths of the radiative recombination continua, and the



Figure 2: *Chandra* LETGS X-ray spectrum of NGC 5548 (Kaastra et al. 2000). The inset shows a zoom of the OVII triplet to which a resonance line, two intercombination lines (unresolved), and a dipole-forbidden line contribute.

strengths of the Fe-L complexes (e.g., Liedahl et al. 1990).

X-ray absorption lines, ionized absorber. With *ROSAT*, the signatures of so-called 'warm' absorbers, absorption edges of highly ionized oxygen ions at $E_{\rm OVII} = 0.74$ keV and $E_{\rm OVIII} = 0.87$ keV, were first detected in MCG−6-30-15 (Nandra & Pounds 1992), following earlier *Einstein* evidence for highly ionized absorbing material in AGN (Halpern 1984). Detailed studies of many other AGN followed, and the signatures of warm absorbers have now been seen in about 50% of the well-studied Seyfert galaxies (see Komossa 1999 for a review). First constraints place the bulk of the ionized material outside the BLR, and depending on its covering factor and location, the warm absorber may be one of the most massive components of the active nucleus. Evidence for ionized absorption was also found in some very high-redshift quasars, starting with observations of PKS 2351-154 (Schartel et al. 1997). Some (but not all) warm absorbers were suggested to contain dust, based on otherwise contradictory optical–X-ray observations (e.g., Brandt et al. 1996, Komossa & Fink 1997b, Komossa & Bade 1998). The first possible detection of Fe-L dust features in the X-ray spectra of MCG−6-30-15 and Mrk 766 was recently reported by Lee et al. (2001) and Lee (2001).

The high-resolution spectrum of the Seyfert galaxy NGC 5548, obtained with the *Chandra* Low Energy Transmission Grating Spectrometer (LETGS), shows many narrow absorption lines of highly ionized metal ions of ogygen,



Figure 3: Soft X-ray spectrum of an AGN, plotted as log flux [a.u.] versus log Energy [keV]. The dashed line shows the input continuum spectrum. The thick solid line gives the spectrum after passage of a warm absorber. The calculation was carried out with the photoionization code *Cloudy* (Ferland 1993). Input parameters (ionization parameter U, column density $N_{\rm w}$) were chosen similar to those obtained from a Beppo-SAX observation of Mrk 766 (Matt et al. 2000) except that U was lowered. Only the absorption edges are shown, and are labeled in the graph (absorption lines were omitted). If this theoretical absorption spectrum is now re-fit, without knowledge of the intrinsic continuum (its shape and level), two fundamentally different solutions are possible: (i) a high-level steep continuum in which case the absorption-solution is recovered, or (ii) a low-level flat continuum (the horizontal thin solid line in the graph) in which case the presence of a strong soft excess (at lower energies than the OVII edge) plus some very broad emission lines are inferred.

neon, iron, etc. (Fig. 2), confirming the presence of a warm absorber in this galaxy (Kaastra et al. 2000). Similar signatures of ionized material have been detected with *Chandra* and *XMM-Newton* in several AGN, including NGC 3783 (Kaspi et al. 2000), IRAS 13349+2438 (Sako et al. 2001), NGC 4051 (Collinge et al. 2001), Mrk 509 (Yaqoob et al. 2002), and MCG −6-30-15 (see next paragraph). First results show that the ionized absorption is complex with a range in ionization states.

Assuming that the ionized absorber outflow is driven by radiation pressure of the central continuum sources, Morales & Fabian (2001) demonstrated that observations can then be used for an estimate of black holes masses in AGN. They derive masses of $M_{\rm BH} \simeq 10^{6.5-7}$ M$_\odot$ for the galaxies of their sample.

Accretion-disk region, Fe-K line. The most direct probe of the black hole region, and particularly, of special and general relativistic effects, is emission from the inner part of the accretion disk (see Fabian 2001 for a review). Tanaka et al. (1995) reported the detection of a broadened FeKα line in MCG$-$6-30-15. The line profile is well explained by the special relativistic effects of beaming and transverse Doppler effect, and the general relativistic effect of gravitational redshift. Depending on details of modeling the continuum, broad-winged Fe lines may also be present in several XMM spectra of AGN (Nandra 2001). At certain times, the red wing of MCG$-$6-30-15 is very broad, extending down to very soft energies (Wilms et al. 2001).

With *XMM* it has also become clearer that the Fe-line profiles are complex, and the line has several sides of formation, including likely the BLR (NGC 5548), the torus (NGC 3783, Mrk 205), the X-ray ionization cone of NGC 1068, and a contribution from the outer parts of the accretion disk (MCG$-$6-30-15).

The case of MCG $-$6-30-15, X-ray spectral complexity. Whereas the signature of an ionized absorber in the form of narrow absorption lines was detected in this galaxy with both, XMM-Newton (Branduardi-Raymont et al. 2001) and *Chandra* (Lee et al. 2001), new interpretations of some of the spectral features were put forward: Branduardi-Raymont et al. suggested that the dominant soft X-ray features, so far interpreted as metal absorption edges of the warm absorber, can be better understood in terms of relativistically broadened emission lines which originate in the accretion disk. On the other hand, Matsumoto & Inoue (2001) noted that the *ASCA*-detected broad wing of the iron K line - so far interpreted in terms of relativistic broadening due to the line's origin in the inner parts of the accretion disk - could be successfully modeled by invoking a two-component warm absorber.

Fig. 3 summarizes and visualizes one of the basic underlying ideas in the discussion of emission versus absorption features at soft X-ray energies (for many additional details see Branduardi-Raymont et al. 2001 and Lee et al. 2001): Depending on where the continuum is placed in Fig. 3, one would either infer the presence of *a huge soft excess plus broad emission lines*, or a powerlaw spectrum modified by *absorption edges*. We do not discuss these ideas further here, except for noting that a disk-line interpretation of the bulk of the soft X-ray features of MCG$-$6-30-15 would leave the puzzle of the discrepant optical and X-ray absorption of this galaxy unanswered, which could be solved by introducing a dusty warm absorber (Reynolds et al. 1997).

Deep high-resolution X-ray observations of, and a search for, variability of the spectral features will be a very important next step in disentangling all components of the complex X-ray spectrum of this galaxy.

1.3 The X-ray search for SMBHs in ULIRGs and LINERs

It is now generally believed that *active* galactic nuclei (AGN) are powered by accretion onto SMBHs. The search for heavily obscured SMBHs in ultraluminous infrared galaxies (ULIRGs), and for low-luminosity AGN (LLAGN) in LINER galaxies is another interesting topic. It will only be briefly touched here, since the emphasis of this review will be on recent evidence for SMBHs in non-active 'normal' galaxies (next Section).

1.3.1 ULIRGs

ULIRGs, characterized by their huge power-output in the infrared which exceeds $10^{12}\,L_\odot$ (Sanders & Mirabel 1996), are powered by massive starbursts or SMBHs. The discussion, which one actually dominates received a lot of attention in recent years (e.g., Joseph 1999, Sanders 1999). In particular, only a small fraction of ULIRGs show AGN signatures in their optical and infrared spectra. Do the remaining ones nevertheless harbor AGN ? X-ray variability and luminous hard X-ray emission are excellent indicators of obscured AGN activity.

With a redshift $z = 0.024$ and a far-infrared luminosity of $\sim 10^{12} L_\odot$, NGC 6240 is one of the nearest members of the class of ULIRGs. Whereas X-rays from distant Hyperluminous IR galaxies, HyLIRGs, were not detected by Wilman et al. (1999), and the ULIRGs in the study of Rigopoulou et al. (1996) were X-ray weak, NGC 6240 turned out to be exceptionally X-ray luminous. Starburst-driven superwinds are the most likely interpretation of the extended emission (see Schulz & Komossa 1999 for alternatives), albeit being pushed to their limits to explain the huge power output (Schulz et al. 1998). The hard spectral component present in the *ROSAT* energy band was interpreted as scattered emission from an obscured AGN (Schulz et al. 1998, Komossa et al. 1998) which shows up more clearly at higher energies, up to 100 keV (e.g., Vignati et al. 1999, Mitsuda 1995, Ikebe et al. 2000). The intrinsically luminous AGN ($L_{\rm x} \approx 10^{44}$ erg/s), can account for at least a substantial fraction of the FIR power output of NGC 6240.

Using *ASCA*, Nakagawa et al. (1999) studied the hard X-ray properties of a sample of 10 ULIRGs. Among these, 50% have hard X-ray detections. The most stringent upper limit for the presence of any hard X-ray emission was reported for Arp 220. The possibility that it is a Compton-thick source cannot be excluded, though.

1.3.2 LINERs

LINER (Low-Ionization Nuclear Emission-Line Region) galaxies are characterized by their optical emission line spectrum which shows a lower degree of ionization than AGN. Their major power source and line excitation mechanism have been a subject of lively debate ever since their discovery. LINERs manifest the most common type of activity in the local universe. If powered by accretion, they probably represent the low-luminosity end of the quasar phenomenon, and their presence has relevance to, e.g., the evolution of quasars, the faint end of the Seyfert luminosity function, the soft X-ray background, and the presence of SMBHs in nearby galaxies.

The X-ray properties of LINERs are inhomogeneous. Spectra of a sample of objects studied by Komossa et al. (1999) are best described by a composition of soft thermal emission and a powerlaw with varying relative contributions of the two components from object to object. Several studies of individual objects are consistent with these results (e.g., Mushotzky 1982, Koratkar et al. 1995, Cui et al. 1997, Ptak et al. 1999, Roberts et al. 1999). X-ray luminosities are in the range $\sim 10^{38-41}$ erg/s; below those typically observed for Seyfert galaxies. The general absence of short-time scale (hours-weeks) X-ray variability (Ptak et al. 1998, Komossa et al. 1999) is consistent with the suggestion that LINERs accrete in the advection-dominated mode (e.g, Yi & Boughn 1998, 1999, and references therein). However, clear positive X-ray detections of LLAGNs in LINERs are still rare. One potential problem problem is to distinguish powerlaw emission of the X-ray binary population of the host galaxy from that of a genuine LLAGN. First *Chandra* results on LINERs show that few, if any, are obscured by absorbers of high column density (Ho et al. 2001). Four out of eight LINERs of that study possess compact nuclear cores, consistent with AGNs.

1.4 The X-ray search for SMBHs in *non-active* ('normal') galaxies, and tidal disruption flares as probes

How can we find *dormant* SMBHs in *non-active* galaxies ? Lidskii & Ozernoi (1979) and Rees (1988) suggested to use the flare of electromagnetic radiation produced when a star is tidally disrupted and accreted by a SMBH as a means to detect SMBHs in nearby *non-active* galaxies.

A star on a near-radial 'loss-cone' orbit gets tidally disrupted once the tidal gravitational forces exerted by the black hole exceed the self-gravitational force of the star (e.g., Hills 1975, Lidskii & Ozernoi 1979, Diener et al. 1997). The tidal radius is given by

$$r_{\rm t} \simeq 7\,10^{12}\,(\frac{M_{\rm BH}}{10^6 M_{\odot}})^{\frac{1}{3}}(\frac{M_*}{M_{\odot}})^{-\frac{1}{3}}\frac{r_*}{r_{\odot}}\ {\rm cm}\,. \qquad (1)$$

The star is first heavily deformed, then disrupted. About 50%–90% of the gaseous debris becomes unbound and is lost from the system (e.g., Young et al. 1977, Ayal et al. 2000). The rest will eventually be accreted by the black hole (e.g., Cannizzo et al. 1990, Loeb & Ulmer 1997). The stellar material, first spread over a number of orbits, quickly circularizes (e.g., Rees 1988, Cannizzo et al. 1990) due to the action of strong shocks when the most tightly bound debris interacts with other parts of the stream (e.g., Kim et al. 1999). Most orbital periods will then be within a few times the period of the most tightly bound matter (e.g., Evans & Kochanek 1989; see also Nolthenius & Katz 1982, Luminet & Marck 1985).

Explicit predictions of the emitted spectrum and luminosity during the disruption process and the start of the accretion phase are still rare (see Sect. 3 for details). The emission is likely peaked in the soft X-ray or UV portion of the spectrum, initially (e.g., Rees 1988, Kim et al. 1999, Cannizzo et al. 1990; see also Sembay & West 1993).

Table 1: Summary of the X-ray and optical properties of the flaring normal galaxies during outburst. z gives the redshift, $T_{\rm bb}$ is the black body temperature derived from a black body fit to the X-ray high-state spectrum (cold absorption was fixed to the Galactic value in the direction of the individual galaxies), 'no emi.' means: no optical emission lines were detected. $L_{\rm x,bb}$ gives the intrinsic luminosity in the (0.1–2.4) keV band, based on the black body fit. This is a lower limit to the actual peak luminosity, since we most likely have not caught the sources exactly at maximum light, since the spectrum may extend into the EUV, and since it was conservatively assumed that no additional X-ray absorption occurs intrinsic to the galaxies.

galaxy name	z	opt. type	$kT_{\rm bb}$ [keV]	$L_{\rm x,bb}$ [erg/s]
NGC 5905	0.011	HII	0.06	$3\ 10^{42*}$
RXJ1242−1119	0.050	no emi.	0.06	$9\ 10^{43}$
RXJ1624+7554	0.064	no emi.	0.097	$\sim 10^{44}$
RXJ1420+5334	0.147	no emi.	0.04	$8\ 10^{43}$
RXJ1331−3243	0.051	no emi.		

[*] Mean luminosity during the outburst; since the flux varied by a factor ∼3 during the observation, the peak luminosity is higher.

2 Giant-amplitude X-ray flares from *non-active* galaxies

2.1 Summary of the original X-ray observations

With the X-ray satellite *ROSAT*, some rather unusual observations have been made in the last few years: the detection of giant-amplitude, non-recurrent X-ray outbursts from a handful of *optically non-active* galaxies, starting with the case of NGC 5905 (Bade et al. 1996, Komossa & Bade 1999). Based on the huge observed outburst luminosity, the observations were interpreted in terms of tidal disruption events. Below, we first give a brief review of the



Figure 4: Optical images of NGC 5905 (right) and RXJ1242−1119 (left). The circles mark the positional uncertainty of the X-ray emission.

properties of all published X-ray flaring non-active galaxies and then discuss the favored outburst scenario. A Hubble constant of $H_0 = 50$ km/s/Mpc is adopted throughout the paper.

So far, four X-ray flaring *non-active* galaxies have been detected: NGC 5905 (Bade et al. 1996, Komossa & Bade 1999; see Fig. 4), RXJ1242−1119 (Komossa & Greiner 1999), RXJ1624+7554 (Grupe et al. 1999), and RXJ1420+5334 (Greiner et al. 2000)[1]; first results on a fifth candidate were presented by Reiprich & Greiner (2001). Based on the position they report, we refer to this source as RXJ1331−3243. All these galaxies show similar properties:

- huge X-ray peak luminosity (up to $\sim 10^{44}$ erg/s),
- giant amplitude of variability (up to a factor ~ 200),
- ultra-soft X-ray spectrum ($kT_{\rm bb} \simeq 0.04$–0.1 keV when a black body model is applied).

A summary of the observations is provided in Table 1. In Fig. 7 X-ray lightcurves of NGC 5905 and RXJ1420+53 are overplotted, shifted in time to the same date of outburst to allow direct comparison. So far, the best sampled lightcurve is that of NGC 5905. The 'merged' lightcurve is consistent with a fast rise and a decline on a time scale of months to years. We performed a preliminary analysis of an archival *ASCA* observation of NGC 5905 carried out in 1999. The flux of NGC 5905 did not drop further compared to the last *ROSAT* observations.

---

[1] The X-ray position error circle of RXJ1420+53 contains a second galaxy for which an optical spectrum is not yet available. Based on the galaxy's morphology, Greiner et al. argue that it is likely non-active.

2.2 Optical observations

Given the unusual X-ray properties of these galaxies, a very important question was: what is the optical classification of the flaring galaxies. Particularly: are there any hints of weak permanent Seyfert activity ? This question is of great interest when discussing outburst scenarios[2].

Optical spectra were taken at different times and at different telescopes, starting several years after the X-ray high-states. The optical spectrum of NGC 5905 turned out to be of HII-type, consistent with its classification prior to the observed X-ray outburst. Neither broad wings in the Balmer lines, nor AGN-typical emission-line ratios, nor high-ionization lines, which usually indicate the presence of an AGN, were detected. The spectra of the other galaxies only show absorption lines from the host galaxies.

Table 2: Coordinates (J 2000) of the optical centers of the galaxies identified as counterparts to the X-ray flares (NGC 5905, RXJ1242−1119A, RXJ1624+7554) or X-ray position (RXJ1420+5334, RXJ1331−3243).

galaxy name	coordinates RA	 DEC
NGC 5905	$15^h15^m23.4^s$	$+55^o31'02''$
RXJ1242−1119A	$12^h42^m38.5^s$	$-11^o19'21''$
RXJ1624+7554	$16^h24^m56.5^s$	$+75^o54'56''$
RXJ1420+5334	$14^h20^m24.2^s$	$+53^o34'11''$
RXJ1331−3243	$13^h31^m57.6^s$	$-32^o43'20''$

2.3 Radio observations

Radio observations are important for two reasons: Firstly, they allow the search for a peculiar, optically hidden AGN at the center of each flaring galaxy (as already discussed by Komossa & Bade (1999) this possibility is a very unlikely explanation for the X-ray flares. It is very important, though, to safely *exclude* exotic AGN scenarios). Besides hard X-ray observations, compact radio emission is a good indicator of AGN activity because radio photons can penetrate even high-column density dusty gas which is not transparent to optical or soft X-ray photons. Secondly, radio emission could possibly be produced in relation to the X-ray flare itself.

---

[2] For instance, theorists working on tidal disruption of stars repeatedly argued that it would be important to be certain about the (optically) *non-active* nature of the flaring galaxy, to exclude AGN-related variability mechanisms (changes in the accretion disk).

2.3.1 NVSS and FIRST search for radio emission from the X-ray outbursters

A search for radio emission from the X-ray flaring galaxies was performed, utilizing the NRAO VLA Sky Survey (NVSS) catalogue (Condon et al. 1998) which contains the results of a 1.4 GHz radio sky survey north of δ=–40°. Except NGC 5905, no flaring galaxy has a NVSS detection. At 1.4 GHz, the emission of NGC 5905 appears extended and is thus related to the galaxy instead of the nucleus (see also next Section).

NGC 5905 is also detected in the FIRST VLA sky survey at 20cm (e.g., Becker et al. 1995). No radio emission from RXJ1420+53 was found. The FIRST catalogue detection limit at the source position is 0.96 mJy/beam. None of the other outbursters is located within a FIRST survey field.

2.3.2 Radio emission from NGC 5905

21 cm neutral hydrogen line emission was detected by van Moorsel (1982; see also Staveley-Smith & Davies 1987), using the Westerbork Synthesis Radio Telescope (WSRT). The emission is spatially resolved (Fig. 3 of van Moorsel 1982) with an extent of diameter 7.3′. Peaks in the HI emission closely follow the spiral arms. Whereas the bulk of the radio emission detected at the frequency of the 21cm line is unrelated to the nucleus, van Moorsel also briefly mentions the presence of unresolved continuum emission of 13.2 mJy.

Extended radio emission was also found by Hummel et al. (1987) at 1.49 GHz, whereas Brosch & Krumm (1984) reported upper limits at 5 GHz (for both, extended emission and a nuclear source). Israel & Mahoney (1990) give an upper limit at 57.5 MHz (see our Fig. 5).

Hummel et al. (1987) reported the presence of an unresolved ($\lesssim 2''$) core source at a frequency of 1.49 GHz. Similar sources were found in 41% of the HII galaxies of the 'complete sample' of Hummel et al. (1987; their Tab. 1).

Finally, we note that the NVSS and FIRST surveys were performed after the X-ray outburst of NGC 5905. However, the NVSS value is consistent with previous measurements of extended radio emission at that frequency. Similarly, the FIRST value is consistent with the pre-flare measurement of Hummel et al. (1987) at the same frequency.

In order to search for a radio source at the nucleus of NGC 5905 after the X-ray outburst (besides a permanent AGN, radio emission could be produced in relation to the X-ray flare) a radio observation with the VLA A array at 8.46 GHz was carried out by M. Dahlem in 1996. No radio source was detected within the central field of view of $100''\times100''$ with a 5σ upper limit for the presence of a central point source of 0.15 mJy (Komossa & Dahlem 2001). Assuming a distance of 75.4 Mpc of NGC 5905 this translates into an upper limit of $L_{8.46\mathrm{GHz}} \leq 1.0\,10^{20}$ W/Hz.

The radio measurements of NGC 5905 are summarized in Fig. 5 together with multi-wavelength observations.



Figure 5: Multi-wavelength continuum spectrum (SED) of NGC 5905 (symbols). The SED of the active galaxy NGC 4151 (Komossa 2001) is shown for comparison by the solid/dotted line. Arrows denote upper limits. Radio data, from left to right: arrow: Clark Lake Radio Observatory TPT array (Israel & Mahoney 1990); lozenge: 21 cm line(van Moorsel 1982); cross: NVSS survey; open circle: FIRST survey; plus: VLA (Hummel et al. 1987); upper limits: WSRT (upper point: total emission within 2.4′, lower point: nuclear emission within 10″; Brosch & Krumm 1984); upper limit: VLA post-flare observation (Komossa & Dahlem 2001); filled squares: IRAS (taken from NED data base), open square: optical B-magnitude (taken from NED), asterisks: X-ray high-state emission (Komossa & Bade 1999). Note: data were taken with different aperture sizes and resolution, and at different times.

2.3.3 Origin of the radio emission of NGC 5905

A large contribution to the radio emission of NGC 5905 comes from the 21cm line of neutral hydrogen. In addition there is some radio continuum emission at the same frequency, and at 1.49 GHz (FIRST). No other radio detection was reported.

The radio emission of large samples of spiral galaxies was examined by, e.g., Brosch & Krumm (1984), Hummel et al. 1987, Giuricin et al. (1990), Israel & Mahoney (1990), Sadler et al. (1995), and Falcke (2001). Radio emission (extended and from the inner few arcseconds) was generally detected from a number of the non-active spiral galaxies in the samples. E.g., Hummel et al. find unresolved ($\lesssim 2''$) core sources in 41% of the HII galaxies of their 'complete sample' at 1.49 GHz. At 57.5 MHz, Israel & Mahoney (1990) detected 68 out of 133 observed galaxies. Trends were repeatedly reported in the literature, that paired HII galaxies (like NGC 5905) and interacting galaxies show enhanced total and central radio emission compared to isolated galaxies (e.g., Giuricin et al. 1990, and references therein). Enhanced star-formation activity was considered a possible explanation of this effect. In some cases, nuclear radio sources in spirals could possibly be explained by radio supernovae (Sadler et al. 1995).

In summary, the radio emission of NGC 5905 is not unusual for its class and does not indicate the presence of a luminous, optically hidden AGN[3].



Figure 6: Locations of the X-ray flaring galaxies on the sky (Galactic II coordinates in aitoff projection).

---

[3] The radio observations do not exclude the presence of some low-level nuclear radio activity related to a low-luminosity AGN, like it may be present for instance in many LINERs (e.g., Falcke 2001, and references therein).



Figure 7: X-ray light curve of NGC 5905 (black squares), and RXJ1420+53 (open circles; shifted in time to match outburst date of NGC 5905). Arrows denote upper limits. A preliminary analysis of an archival *ASCA* observation of NGC 5905 carried out in 1999 shows that the source flux did not drop further compared to the last *ROSAT* observations.

3 Favored outburst scenario: tidal disruption of a star by a SMBH

3.1 (Rejected or unlikely) alternatives to tidal disruption: stellar sources, lensing, GRB, hidden AGN

Most examined outburst scenarios do not survive close scrutiny (Komossa & Bade 1999), because they cannot explain the huge maximum luminosity (e.g., X-ray binaries within the galaxies, or a supernova in a dense medium), are inconsistent with the optical observations (gravitational lensing), or predict a different temporal behavior (X-ray afterglow of a Gamma-ray burst; see, e.g., Fig. 2 of Bradt et al. 2001). Standard AGN scenarios cannot account for the soft X-ray flares and the absence of optical AGN-like emission lines (see the discussion by Komossa & Bade 1999 and Komossa & Voges 2001 for more details).

3.2 Tidal disruption model

Except possibly for some GRB-related emission mechanisms, the huge peak outburst luminosity nearly inevitably calls for the presence of a SMBH[4]. This, in combination with the complete absence of any signs of AGN activity at all wavelengths, makes tidal disruption of a star by a SMBH the favored outburst mechanism.

After a short overview of aspects of tidal disruption models discussed in the literature, we apply these models to the X-ray flare observations.

3.2.1 Tidal disruption, short overview

Historically, tidal disruption of captured stars by black holes was first considered in relation to star clusters and galactic nuclei (e.g., Frank & Rees 1976), and was applied to the nuclei of active galaxies where it was suggested as a means of fueling AGN (e.g., Hills 1975, Sanders 1984), or to explain UV-X-ray variability of AGN (e.g., Kato & Hoshi 1978). Shields & Wheeler (1978) then argued that tidal disruptions of captured stars cannot provide an effective source of fuel of AGN, basically because of two problems: Firstly, the disruption rate is not high enough to sustain a permanent gas flow, and low-angular momentum orbits are quickly depleted of stars. If, however, the stellar density close to the black hole is high enough, stellar collisions would dominate the gas supply over tidal disruption. Secondly, it is difficult to account for the luminosity of the most luminous quasars, since these have masses where the tidal radius is inside the Schwarzschild radius.

A star will only be disrupted if its tidal radius lies outside the Schwarzschild radius of the black hole, else it is swallowed as a whole. This happens for black hole masses larger than $\simeq 10^8$ $M_\odot$; in case of a Kerr black hole, tidal disruption may occur for even larger BH masses if the star approaches from a favorable direction (Beloborodov et al. 1992). More massive black holes may still strip the atmospheres of giant stars.

Due to the complexity of the problem, theoretical work focussed on different subtopics of the complete problem, and on stars of solar mass and radius. Calculations and numerical simulations of the disruption process, the stream-stream collision, the accretion phase, the changes in the stellar distribution of the surroundings and the depopulation and refilling of low-angular momentum orbits, and the disruption rates have been studied in the literature (e.g., Nolthenius & Katz 1982, 1983, Carter & Luminet 1985, Luminet & Marck 1985, Evans & Kochanek 1989, Laguna et al. 1993, Diener et al. 1997, Ayal et al. 2000, Ivanov & Novikov 2001; Kochanek 1994, Lee et al. 1995, Kim et al. 1999; Hills 1975, Gurzadyan & Ozernoi 1979, 1980, Cannizzo et al. 1990, Loeb & Ulmer 1997, Ulmer et al. 1998; Frank & Rees 1976, Lightman & Shapiro 1977, Norman & Silk 1983, Sanders & van Oosterom 1984, Rauch & Ingalls 1998, Rauch 1999; Syer & Ulmer 1999, Magorrian & Tremaine 1999).

---

[4] in fact, even most GRB scenarios involve the presence of a black hole as ultimate energy reservoir

DiStefano et al. (2001) recently considered the case of $M > M_\odot$, and suggested that remnants of tidally stripped stars might be detected as supersoft X-ray sources at the centers of nearby galaxies.



Figure 8: Run of tidal radius and Schwarzschild radius in dependence of black hole mass, for a star of solar mass and radius. For large SMBH masses, stars are swallowed whole, since the tidal radius is no longer outside the Schwarzschild radius.

3.2.2 Tidal disruption in *active* galaxies (?)

During the last several years, tidal disruption was occasionally considered as explanation of some properties of *active* galaxies (either AGN as a class, or individual peculiar observations), although alternative interpretations existed in each case: Tidal disruption was applied by Eracleous et al. (1995) in a duty cycle model to explain the UV brightness/darkness of LINERs. Roos (e.g., 1992) suggested an origin of the BLR clouds of AGN in terms of the gaseous debris of tidally disrupted stars. Peterson & Ferland (1986) proposed tidal disruption as possible explanation for the transient brightening and broadening of the HeII line observed in the Seyfert galaxy NGC 5548. Brandt et al. (1995) and Grupe et al. (1995) reported the detection of an X-ray outburst from the active galaxy Zwicky 159.034 (IC 3599). Besides other outburst mechanisms, tidal disruption was briefly mentioned as possibility. Based on high-resolution post-outburst optical spectra, Komossa & Bade (1999) classified IC3599 as Seyfert type 1.9. In the UV spectral region, two UV spikes were detected at and near the center of the elliptical galaxy NGC 4552. The central flare, although rather weak, was interpreted by Renzini et al. (1995) as accretion event (the tidal stripping of a star's atmosphere by a SMBH, or the accretion of a molecular cloud). There are several indications (e.g., from radio observations), that NGC 4552 shows permanent low-level activity.

3.2.3 Emission of radiation, temporal evolution, and model uncertainties

Intense electromagnetic radiation will be emitted in three phases of the disruption and accretion process: First, during the stream-stream collision when different parts of the bound stellar debris first interact with themselves (Rees 1988). Kim et al. (1999) have carried out numerical simulations of this process and find that the initial burst due to the collision may reach a luminosity of 10^{41} erg/s, under the assumption of a BH mass of 10^6 $M_\odot$ and a star of solar mass and radius. Secondly, radiation is emitted during the accretion of the stellar gaseous debris. Finally, the unbound stellar material leaving the system may shock the surrounding interstellar matter and cause intense emission, like in a supernova remnant (Khokhlov & Melia 1996).

Many details of the tidal disruption and the related processes are still unclear. In particular, the flares cannot be standardised. Observations would depend on many parameters, like the type of disrupted star, the impact parameter, the spin of the black hole, effects of relativistic precession, and the complication of radiative transfer by effects of viscosity and shocks (Rees 1990). Uncertainties also include the amount of the stellar material that is accreted (part may be ejected as a thick wind, or swallowed immediately). Related to this is the duration of the flare-like activity, which may be months or years to tens of years (e.g., Rees 1988, Cannizzo et al. 1990, Gurzadyan & Ozernoi 1979), followed by a decline on a longer time scale. The flare duration depends on how fast the stellar debris circularizes and how fast it accretes. If both time scales are short and the material accretes at the same rate as it falls back towards the black hole, the decline time scale of the flare scales as

$$\frac{dm}{dt} \propto t^{-\frac{5}{3}} \tag{2}$$

(e.g., Evans & Kochanek 1989, Rees 1990). Else energy output will be spread more evenly in time (Cannizzo et al. 1990).

Another uncertainty in predicting the flare spectrum arises from the mode of accretion: does it proceed via a thin disk (e.g., Cannizzo et al. 1990) or a thick disk (Ulmer et al. 1998), and, if in a thick disk, under which angle do we view it ?

3.3 Order of magnitude estimates and consistency checks

3.3.1 Inferences from X-ray observations

Although many details of the actual tidal disruption process are still unclear, some basic predictions have been repeatedly made in the literature how a tidal disruption event should manifest itself observationally:

- (1) the event should be of finite duration (a 'flare'),
- (2) it should be very luminous (up to $L_{\max} \approx 10^{45}$ erg/s in maximum), and

- (3) it should reside in a galaxy which is otherwise perfectly *non*-active (to be sure to exclude an upward fluctuation in gaseous accretion rate of an *active* galaxy).

All three predictions are fulfilled by the X-ray flaring galaxies; particularly by NGC 5905 and RXJ1242−1119, which are the two best-studied cases so far.

In addition, we can do some further order of magnitude estimates and consistency checks. The luminosity emitted if the black hole is accreting at its Eddington luminosity can be estimated by

$$L_{\rm edd} = \frac{4\pi G M m_{\rm p} c}{\sigma_{\rm T}} \simeq 1.3\times10^{38} M/M_{\odot}\ {\rm erg/s}\,. \qquad (3)$$

In case of NGC 5905, a BH mass of at least a few $\sim 10^4$ $M_{\odot}$ would be required to produce the observed luminosity. This is a *lower limit* on the black hole mass, since we likely did not observe $L_{\rm x}$ at its peak value due to observational gaps in the lightcurve, and since other conservative assumptions were made[5]. A comparison with the SMBH mass of NGC 5905 using indirect *optical* methods is performed in Sect. 3.3.2. For the other galaxies, using again $L_{\rm edd}$, we infer BH masses reaching up to a few 10^6 $M_{\odot}$. This is, again, a lower limit. Alternative to a complete disruption event, the atmosphere of a giant star could have been stripped.

In a simple black body approximation, the temperature of the accretion disk scales with black hole mass as

$$T \simeq 8\,10^4\,(\frac{M_{\rm BH}}{M_{\odot}})^{\frac{1}{12}}\ {\rm K}\ \ ({\rm at}\ r_t), \quad T \simeq 2\,10^7\,(\frac{M_{\rm BH}}{M_{\odot}})^{-\frac{1}{4}}\ {\rm K}\ \ ({\rm at}\ 3\,r_S)\,. \qquad (4)$$

This gives $T_{\rm r_{tidal}} \simeq 3\,10^5$ K, $T_{\rm 3r_S} \simeq 7\,10^5$ K for M=10^6 $M_{\odot}$, where $r_{\rm S}$ is the Schwarzschild radius. Using black body fits of the X-ray flare spectra we find temperatures in a similar range; $T_{\rm obs} \simeq$ (4-10) 10^5 K. Like in AGN, X-ray powerlaw tails could develop. They might have escaped detection during the observations, since weak, or they may develop only after a certain time after the start of the accretion phase. We soon expect first results from a *Chandra* and *XMM* observation of RXJ1242−1119, which will give valuable constraints on the post-flare evolution.

The Eddington time scale for the accretion of the stellar material is given by

$$t_{\rm edd} \simeq 4\,\eta_{0.1}(M_{\rm BH}/10^6 M_{\odot})(M_*/0.1 M_{\odot})\ {\rm yrs}\,. \qquad (5)$$

Uncertainties in estimating the total duration of the tidal disruption event arise from questions like: how much material is actually accreted or expelled, does a strong wind develop, etc. (see Sect. 3.2.3). The events are expected to last for months to years (e.g., Rees 1988). Observationally, the duration of the events was at least several days, followed by gaps in the observations. The source fluxes were then significantly down several years later (e.g., Fig. 9 of Komossa & Bade 1999). Apart from theoretical uncertainties in the prediction of the decline time scale in the total luminosity output (Sect. 3.2.3), the emission will also likely shift outside the *ROSAT* band, from the X-ray to the EUV-UV band, with increasing time.

---

[5] e.g., the amount of absorption was fixed to the lowest possible value, $N_{\rm Gal}$

Finally, we note that the redshift distribution of the few sources observed so far is consistent with the predicted tidal disruption rate, in the sense that the events are sufficiently distant to define a large volume of space, in which the detection of a few events would be expected.

3.3.2 Non X-ray estimates of the black hole mass in NGC 5905

An optical rotation curve was obtained by Komossa & Bade (1999) which allowed an estimate of the mass enclosed within 0.7 kpc: $M \approx 10^{10}$ M$_\odot$. This immediately provides an upper limit on BH mass, but the volume sampled is still too large to estimate the actual BH mass. An HST-based rotation curve would significantly improve the above limit and thus the constraints on a central dark mass.

In order to get a better (non-X-ray) estimate of the BH mass of NGC 5905, we used the correlation between bulge properties and BH mass.

NGC 5905 has a total blue magnitude of $\mathrm{m}_{B,0} = 12.1^{\mathrm{m}}$. Using the bulge-to-disk luminosity ratio generally valid for galaxies of the Hubble-type of NGC 5905 (SBb), $k = 0.25$ (Salucci et al. 2000), gives the absolute bulge blue magnitude, $\mathrm{B}_{\mathrm{T,0}}^{\mathrm{bulge}} = -20.5$. We then compared with two recent studies that correlate BH mass and bulge blue luminosity: (i) the work of Ferrarese & Merritt (2000; FM00) which concentrates mostly on elliptical galaxies, and (ii) the work of Salucci et al. (2000; S00) on spiral galaxies.

Using the relation between $\mathrm{B}_{\mathrm{T,0}}^{\mathrm{bulge}}$ and M_{BH} of FM00 (their 'sample A', their Tab. 2; see also, e.g., Franceschini et al. 1998) gives a BH mass of a few times 10^8 M$_\odot$ for NGC 5905. This is close to the limiting BH mass for tidal disruption of a solar-type star to work. However, it has to be kept in mind, that the $\mathrm{B}_{\mathrm{T,0}}^{\mathrm{bulge}}$ - M_{BH} relation shows a very large scatter (in contrast to the M_{BH} - σ relation), and that the few spirals in the sample of FM01 tend to be located below the relation followed by ellipticals. Therefore, in a second step, we used the results of S00 on BH masses in (late-type) spiral galaxies (their Fig. 6). In that case we obtain an upper limit for the BH mass of NGC 5905 of $M_{\mathrm{BH}} \lesssim 10^7$ M$_\odot$.

Finally, the relation between radio luminosity and BH mass of Franceschini et al. (1998, their Fig. 3; see also Wu & Han 2000) was employed. The measured radio upper limit for the nucleus of NGC 5905 then translates into an upper limit on black hole mass of $M_{\mathrm{BH}} \lesssim 2.5\,10^8$ M$_\odot$. Results are summarized in Tab. 3.

3.3.3 Properties of the host galaxies

The expected rate of tidal disruption events is about one event in *at least* $\sim 10^4$ years per galaxy (e.g., Magorrian & Tremaine 1999), and the whole

Table 3: Summary of mass estimates of the black hole at the center of NGC 5905, employing different methods (see the text for details).

energy band	method	BH mass
X-rays	$L_{\rm Eddington}$	$>$ few 10^4 $M_\odot$
optical	$M_{\rm B,bulge} - M_{\rm BH}$ correlation, spirals	$\approx 10^7$ $M_\odot$
	$M_{\rm B,bulge} - M_{\rm BH}$ correlation, ellipticals	$\approx$ few 10^8 $M_\odot$
	rotation curve	$\ll 10^{10}$ $M_\odot$
radio	nuclear radio power $-$ $M_{\rm BH}$ correlation	$< 2.5\,10^8$ $M_\odot$



Figure 9: Position of NGC 5905 in the $L_{\rm x} - L_{\rm B}$ diagram in outburst and low-state. The dashed lines mark the region populated by some samples of elliptical galaxies.

ROSAT data base has to be employed for a systematic search for further tidal disruption events (for first results see Komossa & Bade 1999, and below).

The disruption rate depends on the efficiency with which the loss-cone orbits are re-filled (Frank and Rees 1976, Lightman and Shapiro 1977, Shields & Wheeler 1978). This can be done by perturbations of the stellar orbits, e.g. by merging events (Roos 1981) or a triaxial gravitational potential in the galaxy's core (Norman & Silk 1983).

Are the observed flaring galaxies special in this context (i.e., did any process aid in re-populating the loss-cone orbits)? So far, not much is known about the host galaxies of the few flaring galaxies. The best studied case is NGC 5905. It is interesting to note that this galaxy posses multiple triaxial structures with a secondary bar (Friedli et al. 1996, Wozniak et al. 1995) which might aid occasional tidal disruption events by disturbing the stellar velocity fields. NGC 5905 is in a pair with NGC 5908.

RXJ1242−11 is actually a pair of galaxies at similar redshift, and it is well possible that both galaxies are interacting. The X-ray error circle of another outburster, RXJ1420+53, also includes two galaxies. However, a redshift is so far only available for the brighter of the two.

The Hubble types of the flaring galaxies are not known in all cases. NGC 5905, of type SB (Keenan 1937), is one of the largest spiral galaxies known (e.g., Romanishin 1983), whereas some of the other host galaxies look like ellipticals. Deeper optical imaging is presently in progress.

4 Search for further X-ray flares

While we wait for the next-generation of X-ray all-sky surveys, we can still make use of an existing data base which has not yet been fully exploited, the *ROSAT* data base (Voges et al. 1999).

In a first step to search for further cases of strong X-ray variability the sample of nearby galaxies of Ho et al. (1995) and *ROSAT* all-sky survey (RASS) and archived pointed observations were used. The sample of Ho et al. has the advantage of the availability of optical spectra of good quality, which are necessary when searching for 'truly' non-active galaxies. 136 out of the 486 galaxies in the catalogue were detected in pointed observations. The source countrates were then compared with those measured during the RASS.

4.1 Non-active galaxies

No other X-ray flaring, optically in-active galaxy was found. The absence of any further flaring event among the sample galaxies is entirely consistent with the expected tidal disruption rate of one event in $\sim 10^{4-5}$ years per galaxy (e.g., Magorrian & Tremaine 1999).

The next step, presently in progress, will be an extended search for X-ray flaring events based on the whole *ROSAT* all-sky survey database. This approach will allow statistical inferences on the abundance of SMBHs in non-active galaxies (Sembay & West 1993).

4.2 AGN

Several of the sample galaxies show variability by a factor 10–30. All of these are well-known AGN.

Many active galactic nuclei are variable in X-rays with a range of amplitudes, typically a factor 2–3 (e.g., Mushotzky et al. 1993, Ulrich et al. 1997). The cause of variability is usually linked in one way or another to the central engine; for instance by changes in the accretion disk (e.g., Piro et al. 1988, 1997), or by variable obscuration (e.g., Komossa & Fink 1997a, Komossa & Janek 2000). None of the well-studied X-ray variable AGN of the present sample are candidates for tidal disruption events since their lightcurves show recurrent variability on different time scales.

An example for a highly variable AGN among the present sample galaxies is NGC 4051. Its long-term *ROSAT* X-ray lightcurve exhibits variability in countrate by a factor ∼30. Only a small part of the variability of NGC 4051 can be explained with a variable warm absorber, the rest is likely intrinsic (Komossa & Fink 1997a, Komossa & Janek 2001, and references therein).

Even higher total amplitude of variability is detected in two subsequent *ROSAT* observations of NGC 3516. The X-ray countrate varies by a factor ∼50 (Komossa & Bade 1999, Komossa & Halpern 2001, in prep.); variable cold absorption likely plays a major part in explaining the observations.

5 Future perspectives

X-ray outbursts from non-active galaxies provide important information on the presence of SMBHs in these galaxies, and the link between active and normal galaxies. One advantage of this method compared to other approaches to search for central dark masses – like HST-based galaxy rotation curves – is that the X-ray flare emission originates from the *very vicinity* of the SMBH. Therefore, potentially it provides much tighter constraints on the black hole mass. Flares can also be detected out to larger cosmological distances.

Follow-up studies: future X-ray surveys. Future X-ray surveys, like those planned with the *LOBSTER* ISS X-ray all-sky monitor (Fraser 2001), *MAXI* (Yuan et al. 2001, Mihara 2001), and *ROSITA* (Predehl 2001) will be valuable in finding more of these outstanding sources. In addition, a number of flare events are expected to be detected (Yuan et al. 2002) in pointings of the *XMM-Newton* and *Chandra* missions.

On the one hand, all-sky surveys will be important in detecting the brightest events, due to their large areal sky coverage. Those surveys with sensitivity at the softest energies will be most efficient. On the other hand, deeper pointings on limited areas, or 'pencil beam' surveys will increase the number of (more distant, on average fainter) events (see Yuan et al. 2002 for logN-logS estimates of the expected number of flares detectable with *XMM-Newton*, and Sembay & West 1993 for a general discussion). Concerning deep surveys in limited fields of view, two effects are important. Firstly, there is a limiting maximal brightness the events can reach in the context of the tidal disruption scenario: the most luminous events are those with black hole masses around 10^8 $M_\odot$, with high accretion rate, and the maximal possible fraction emitted in the X-ray band (definite upper limit: $L_{\rm flare} \lesssim 10^{45-46}$ erg/s). Secondly, there is a limiting distance out to which flare events are detectable because the flare spectra are very soft, and for increasing redshift more and more of the (black-body-like) emission is shifted out of the observable energy band (Fig. 10). In addition, distant galaxies may show a higher intrinsic fraction of cold gas which will heavily absorb at soft X-rays.

After the discovery of new X-ray flare events, *rapid follow-up* multi-wavelength observations will be essential. Apart from valuable new constraints on



Figure 10: Simulation of an XMM spectrum (EPIC pn instrument) of an X-ray flare similar to the one observed from NGC 5905, assuming a medium-deep exposure of 100 ksec and a single black body spectrum of $kT = 0.05$ keV absorbed by $N_{\rm Gal} = 1.5\,10^{20}$ cm^{-2} at a redshift of z=0.01. At z=0.1 the very soft black-body-like emission is partly redshifted out of the instrument's energy bandpass, causing a further drop in the observed countrate (∼30% at z=0.1) in addition to the flux decrease because of the larger distance. The effect increases with redshift such that very distant flares can no longer be detected since shifted out of the observable energy range. Among the X-ray flares, that of NGC 5905 was of lowest luminosity. More luminous flares would therefore be detectable out to much higher redshifts. In addition to the continuum spectrum, we may expect to detect emission features which arise in the accreted material, or absorption features from the ISM of the host galaxy.

the favored outburst scenario which would also allow a refinement of model calculations, these observations would enable us to address a number of important topics:

Absorption-line spectroscopy of the IGM/ISM. As the flare emission travels through the ISM of the host galaxy, and the IGM, absorption features will be imprinted on the X-ray spectrum. These can then be used to study the properties of the absorbing material.

Emission-line spectroscopy of the circumnuclear material. If the soft X-ray flare emission has an extension into the EUV, which is highly likely, then optical observations will be important in order to detect potential emission lines that were excited by the outburst emission. Firstly, any gaseous material close to the nucleus is expected to show an emission-line response. The time variability of these lines will allow a reverberation mapping of the circumnuclear gas; line profiles, and line-ratios will allow to estimate the velocity structure and physical conditions (density, abundances) of this gas. In particular, this would also enable us to search for the presence of a BLR in these optically in-active galaxies.

Probing the realm of strong gravity. If observed with high spectral and temporal resolution with the next generation of X-ray telescopes, like the *XEUS* mission sensitive between 0.05–30 keV, the flare spectra may allow to probe the realm of strong gravity, since the temporal evolution of the stellar debris, and of potential spectral features, will depend on relativistic precession effects around the Kerr metric.

Acknowledgements: It is a pleasure to thank the Astronomische Gesellschaft for awarding the Ludwig-Biermann price and for inviting me to give this talk. I would like to thank Weimin Yuan, Michael Dahlem, Jules Halpern, Norbert Bade, Andrew Ulmer, Niel Brandt, Martin Elvis, David Meier, and the participants of the CAS-MPG Workshop on High-Energy Astrophysics at Ringberg Castle for very useful discussions and comments on the subject of X-ray flares from non-active galaxies, and Norbert Bade for taking the optical image of NGC 5905 which is shown in Fig. 4. I gratefully remember Henner Fink for introducing me to the work with X-ray data, for discussions and helpful advice. Henner Fink passed away in December 1996. The *ROSAT* project was supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF/DLR) and the Max-Planck-Society.
Preprints of this and related papers can be retrieved at
http://www.xray.mpe.mpg.de/∼skomossa/

References

Antonucci R., 1993, ARA&A 31, 473

Ayal S., Livio M., Piran T., 2000, ApJ 545, 772

Bade N., Komossa S., Dahlem M., 1996, A&A 309, L35

Becker R.H., White R.L., Helfand D.J., 1995, ApJ, 450, 559

Beloborodov A.M., Illarionov A.F., Ivanov P.B., Polnarev A.G., 1992, MNRAS 259, 209

Bradt H., Levine A.M., Marshall F.E., et al. 2001, to appear in ESO Astrophysics Symposia, F. Frontera et al. (eds), [astro-ph/0108004]

Brandt W.N., Pounds K.A., Fink H.H., 1995, MNRAS 273, L47

Brandt W.N., Fabian A.C., Pounds K.A., 1996, MNRAS 278, 326

Branduardi-Raymont G., Sako M., Kahn S., et al., 2001, A&A 365, L140

Brosch N., Krumm N., 1984, A&A 132, 80

Brown B.A., Bregman J.N., 1998, ApJ 495, L75

Canizares C.R., Fabbiano G., Trinchieri G., 1987, ApJ 312, 503

Cannizzo J.K., Lee H.M., Goodman J., 1990, ApJ 351, 38

Carter B., Luminet J.P., 1985, MNRAS 212, 23

Collin S., Abrassart A., Czerny B., Dumont A.-M., Mouchet M., EDPS Conf. Series in Astron. & Astrophys., in press [astro-ph/0003108]

Collin-Souffrin S., Czerny B., Dumont A.-M., Zycki P.T., 1996, A&A 314, 393

Collinge M.J., Brandt W.N., Kaspi, S., et al., 2001, ApJ 557, 2

Condon J.J., Cotton W.D., Greissen E.W., et al., 1998, AJ 115, 1693

Cui W., Feldkun D., Braun R., 1997, ApJ 477, 693

Diener P., Frolov V.P., Khokhlov A.M., Novikov I.D., Pethick C.J., 1997, ApJ 479, 164

Di Stefano R., Greiner R., Murray S., Garcia M., 2001, ApJ 551, L37

Elvis M., 2000, ApJ 545, 63

Elvis M., 2001, in press, [astro-ph/0109513]

Elvis M., Briel U., Henry J.P., 1983, ApJ 268, 105

Elvis M., Fassnacht C., Wilson A.S., Briel U., 1990, ApJ 361, 459

Eracleous M., Livio M., Binette L., 1995, ApJ 445, L1

Evans C.R., Kochanek C.S., 1989, ApJ 346, L13

Fabian A., 2001, in: TEXAS Symposium, in press [astro-ph/0103438]

Falcke H., 2001, in: Reviews in Modern Astronomy 14, R. Schielicke (ed.), 15

Ferland G.J., 1993, University of Kentucky, Physics Department, Internal Report

Ferrarese L., Merritt D., 2000, ApJ, 539, L9

Ferrarese L., Pogge R.W., Peterson B.M., et al., 2001, ApJ 555, L79

Franceschini A., Vercellone S., Fabian A.C., 1998, MNRAS 297, 817

Frank J., Rees M.J., 1976, MNRAS 176, 633

Fraser G., 2001, in: MAXI workshop on AGN variability, in press

Friedli D., Wozniak H., Rieke M., Martinet L., Bratschi P., 1996, A&AS 118, 461

Gabriel A.H., Jordan C., 1969, MNRAS 145, 241

George I.M., Fabian A.C., 1991, MNRAS 249, 352

Giuricin G., Bertotti G., Mardirossian F., Mezzetti M., 1990, MNRAS 247, 444

Greenhill L.J., Henkel C., Becker R., Wilson T.L., Wouterloot J.G.A., 1995, A&A 304, 21

Greiner J., Schwarz R., Zharikov S., Orio M., 2000, A&A 362, L25

Grupe D., Beuermann K., Mannheim K., et al., 1995, A&A 299, L5

Grupe D., Leighly K., Thomas H., 1999, A&A 351, L30

Gurzadyan V.G., Ozernoi L.M., 1979, Nature 280, 214

Gurzadyan V.G., Ozernoi L.M., 1980, A&A 86, 315

Hills J.G., 1975, Nature 254, 295

Halpern J.P., 1984, ApJ 281, 90

Ho L.C., Filippenko A.V., Sargent W.L.W., 1995, ApJS 98, 477

Ho L.C., et al., 2001, ApJ, in press

Hummel E., van der Hulst J.M., Keel W.C., Kennicutt R.C. Jr., 1987, A&AS 70, 517

Ikebe Y., Leighly K., Tanaka Y., et al., 2000, MNRAS 316, 433

Israel F.P., Mahoney M.J., 1990, ApJ 352, 30

Ivanov P.B., Novikov I.D., 2001, ApJ 549, 467

Joseph R.D., 1999, Ap&SS 266, 321

Kaastra J., Mewe R., Liedahl D.A., Komossa S., Brinkman A.C., 2000, A&A 354, L83

Kaspi S., Brandt W.N., Netzer H., et al., 2001, ApJ 554, 216

Kato M., Hoshi R., 1978, Prog. Theor. Phys. 60/6, 1692

Keenan P.C., 1937, ApJ 85, 325

Khokhlov A., Melia F., 1996, ApJ 457, L61

Kim S.S., Park M.-G., Lee H.M., 1999, ApJ 519, 647

Kochanek C., 1994, ApJ 422, 508

Komossa S., 1999, in: ASCA/ROSAT Workshop on AGN and the X-ray Background, T. Takahashi, H. Inoue (eds), ISAS Report, p. 149; [also available at astro-ph/0001263]

Komossa S., 2001, A&A 371, 507

Komossa S., 2001, in: IX. Marcel Grossmann Meeting on General Relativity, Gravitation and Relativistic Field Theories, V. Gurzadyan et al. (eds), World Scientific, in press

Komossa S., Bade N., 1998, A&A 331, L49

Komossa S., Bade N., 1999, A&A 343, 775

Komossa S., Böhringer H., Huchra J., 1999, A&A 349, 88

Komossa S., Dahlem M., 2001, in: MAXI workshop on AGN variability, ISAS Report, in press

Komossa S., Fink H., 1997a, A&A 322, 719

Komossa S., Fink H., 1997b, A&A 327, 483

Komossa S., Greiner J., 1999, A&A 349, L45

Komossa S., Janek M., 2000, A&A 354, 411

Komossa S., Schulz H., Greiner J., 1998, A&A 334, 110

Komossa S., Voges W., 2001, in: MPG-CAS workshop on high-energy astrophysics, preprint available at: http://www.xray.mpe.mpg.de∼skomossa/publrev.html

Koratkar A., Deustua S.E., Heckman T., et al., 1995, ApJ 440, 132

Kormendy J., Richstone D.O., 1995, ARA&A 33, 581

Kormendy J., Gebhardt K., 2001, in press [astro-ph/0105230]

Krolik J.H., Kriss G.A., 1995, ApJ 447, 512

Laguna P., Miller W.A., Zurek W.H., Davies M.B., 1993, ApJ 410, L83

Lee H.M., Kang H., Ryu D., 1995, ApJ 464, 131

Lee J., Ogle P., Canizares C., et al., 2001, ApJ 554, L13

Lee J., 2001, talk given at: Workshop on X-ray spectroscopy of active galactic nuclei with Chandra and XMM-Newton (Garching, Dec. 2001)

Lidskii V.V., Ozernoi L.M., 1979, Sov. Astron. Lett. 5(1), 16

Liedahl D.A., Kahn S.M., Osterheld A.L., Goldstein W.H., 1990, ApJ 350, L37

Lightman A.P., Shapiro S.L., 1977, ApJ 211, 244

Loeb A., Ulmer A., 1997, ApJ 489, 573

Luminet J.P., Marck J.-A., 1985, MNRAS 212, 57

Lynden-Bell D., 1969, Nature 223, 690

Magorrian J., Tremaine S., 1999, MNRAS 309, 447

Matsumoto C., Inoue H., 2001, in: MAXI workshop on AGN variability, in press

Matt G., Perola G.C., Fiore F., et al., 2000, A&A 363, 863

Mihara T., 2001, in: MAXI workshop on AGN variability, in press

Mitsuda K., 1995, Ann.N.Y.Acad.Sc. 759, Proc. 17th Texas Symp. Relat. Ap. and Cosm., H. Böhringer, G.E. Morfill, J.E. Trümper (eds), 213

Miyoshi M., Moran J., Herrnstein J., et al., 1995, Nature 373, 127

Morales R., Fabian A.C., 2001, MNRAS, in press [astro-ph/0109050]

Mushotzky R., 1982, ApJ 256, 92

Mushotzky R.F., Done C., Pounds K.A., 1993, ARA&A 31, 717

Nakagawa T., Kii T., Fujimoto R., et al., 1999, Ap&SS 266, 43

Nandra P., 2001, talk given at: Workshop on X-ray spectroscopy of active galactic nuclei with Chandra and XMM-Newton (Garching, Dec. 2001)

Nandra K., Pounds K.A., 1992, Nature 359, 215

Netzer H., 1993, ApJ 411, 594

Neufeld D.A., Maloney P.R., 1995, ApJ 447, L17

Nolthenius R.A., Katz J.I, 1982, ApJ 263, 377

Nolthenius R.A., Katz J.I, 1983, ApJ 269, 297

Norman C., Silk J., 1983, ApJ 266, 502

Ogle P.M., Marshall H.L., Lee J.C., Canizares C.R., 2000, ApJ 545, L81

Peterson B.M., Ferland G.J., 1986, Nature 324, 345

Peterson B.M., 2001, in: Advanced lectures on the starburst-AGN connection, I. Aretxaga et al. (eds), World Scientific, 3

Peterson B.M., Wandel A., 2000, ApJ 540, L13

Piro L., Massaro E., Perola G.C., Molteni D., 1988, ApJ 325, L25

Piro L., Balucinska-Church M., Fink H., et al., 1997, A&A 319, 74

Predehl P., 2001, in: MAXI workshop on AGN variability, in press

Ptak A., Yaqoob T., Mushotzky R., Serlemitsos P., Griffiths R., 1998, 501, L37

Ptak A., Serlemitsos P., Yaqoob T., Mushotzky R., 1999, ApJS 120, 179

Rauch K.P., 1999, ApJ 514, 725

Rauch K.P., Ingalls B., 1998, MNRAS 299, 1231

Rees M.J., 1988, Nature 333, 523

Rees M.J., 1989, Rev. mod. Astr. 2, R. Schielicke (ed.), 1

Rees M.J., 1990, Science 247, 817

Rees M.J., 1997, Rev. mod. Astr. 10, R. Schielicke (ed.), 179

Reiprich T., Greiner J., 2001, in: ESO workshop on black holes in binaries and AGN, p. 168

Renzini A., Greggio L., Di Serego Alighieri S., et al., 1995, Nature 378, 39

Reynolds C.S., Ward M.J., Fabian A.C., Celotti A., 1997, MNRAS 291, 493

Rigopoulou D., Lawrence A., Rowan-Robinson M., 1996, MNRAS 278, 1049

Roberts T.P., Warwick R.S., Ohashi T., 1999, MNRAS 304, 52

Romanishin W., 1983, MNRAS 204, 909

Roos N., 1981, A&A 104, 218

Roos N., 1992, ApJ 385, 108

Sadler E., Slee O.B., Reynolds J.E., Roy A.L., 1995, MNRAS 276, 1373

Sako M., Kahn S.M., Behar E., et al., 2001, A&A 365, L168

Sanders D.B., 1999, Ap&SS 266, 331

Sanders D.B., Mirabel I.F., 1996, ARA&A 34, 749

Sanders R.H., 1984, A&A 140, 52

Sanders R.H., van Oosterom W., 1984, A&A 131, 267

Salucci P., Ratnam C., Monaco P., Danese L., 2000, MNRAS 317, 488

Schartel N., Komossa S., Brinkmann W., Fink H.H., Trümper J., Wamsteker W., 1997, A&A 320, 421

Schulz H., Komossa S., 1999, contribution to: Bad Honnef workshop on Trends in Astrophysics and Cosmology, astro-ph/9905118

Schulz H., Komossa S., Berghöfer T., Boer B., 1998, A&A 330, 823

Sembay S., West R.G., 1993, MNRAS 262, 141

Shields G.A., Wheeler J.C., 1978, ApJ 222, 667

Staveley-Smith L., Davies R.D., 1987, MNRAS 224, 953

Svensson R., 1994, ApJS 92, 585

Syer D., Ulmer A., 1999, MNRAS 306, 35

Tanaka Y., Nandra K., Fabian A.C., et al., 1995, Nature 375, 659

Ulmer A., Paczynski B., Goodman J., 1998, A&A 333, 379

Ulrich, M.-H., Maraschi L., Urry C.M., 1997, ARA&A 35, 445

van Moorsel G.A., 1982, A&A 107, 66

Vignati P., Molendi S., Matt G., et al., 1999, A&A 349, L57

Voges W., et al., 1999, A&A 349, 389

Wilman R.J., 1999, Ap&SS 266, 55

Wilms J., Reynolds C., Begelman M.C., et al., 2001, MNRAS L, in press [astro-ph/0110520]

Wozniak H., Friedli D., Martinet L., Martin P., Bratschi P., 1995, ApJS 111, 115

Wu X.-B., Han J.L., 2001, A&A 380, 31

Yaqoob T., George I.M., Turner T.J., 2002, ASP conf. series, in press [astro-ph/0111428]

Yi I., Boughn S.P., 1998, ApJ 499, 198

Yi I., Boughn S.P., 1999, ApJ 515, 579

Young P., Shields G., Wheeler J.C., 1977, ApJ 212, 367

Yuan W., Matsuoka M., Shirasaki Y., et al., 2001, in: MAXI workshop on AGN variability, in press

Yuan W., et al., 2002, in: New Visions of the X-ray Universe, F. Jansen et al. (eds.), in press